

Stroke

American Stroke
AssociationSM

JOURNAL OF THE AMERICAN HEART ASSOCIATION

A Division of American
Heart Association



**Statins After Ischemic Stroke and Transient Ischemic Attack: An Advisory
Statement From the Stroke Council, American Heart Association and American
Stroke Association**

The Stroke Council

Stroke 2004;35;1023-

DOI: 10.1161/01.STR.0000122762.96972.DD

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214
Copyright © 2004 American Heart Association. All rights reserved. Print ISSN: 0039-2499. Online
ISSN: 1524-4628

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://stroke.ahajournals.org/cgi/content/full/35/4/1023>

Subscriptions: Information about subscribing to *Stroke* is online at
<http://stroke.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, 351 West Camden
Street, Baltimore, MD 21202-2436. Phone 410-5280-4050. Fax: 410-528-8550. Email:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/static/html/reprints.html>

Statins After Ischemic Stroke and Transient Ischemic Attack

An Advisory Statement From the Stroke Council, American Heart Association and American Stroke Association

The Stroke Council

Based on results of numerous large-scale randomized trials, the vast majority of patients with a history of ischemic stroke or transient ischemic attack could benefit from statin use.

Although prevention of second stroke was not the primary aim of any completed study, some studies included subjects whose primary reason for entry was stroke. Multiple studies have shown that statins reduce risk of stroke in those with coronary artery disease and elevated total or low-density lipoprotein (LDL) cholesterol. Recently, the Heart Protection Study showed that simvastatin 40 mg/day reduced the risk of stroke by 25% among patients with coronary artery disease, other occlusive arterial disease, or diabetes.¹ In the subgroup enrolled with prior ischemic stroke or transient ischemic attack but no coronary artery disease, the risk of major vascular events (coronary events, stroke, or revascularization) was reduced by 21% (absolute risk reduction, 1% per year; number needed to treat 102 to prevent 1 event each year). Benefits persisted in those with LDL <116 mg/dL or total cholesterol <193 mg/dL. A meta-analysis also shows that the benefits of statins in reducing the rates of stroke and cardio-

vascular events is independent of cholesterol levels and occur with other statins.² Given early benefits in trials of acute coronary syndromes, statin initiation during hospitalization for first ischemic stroke of atherosclerotic origin is probably justified and may increase rates of long-term use. Results of the ongoing SPARCL trial³ will provide additional information about the role of statins in the minority of patients with prior stroke but no history of coronary heart disease, other occlusive arterial disease, or diabetes.

References

1. Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet*. 2002;360:7–22.
2. Corvol JC, Bouzamondo A, Sirol M, Hulot JS, Sanchez P, Lechat P. Differential effects of lipid-lowering therapies on stroke prevention: a meta-analysis of randomized trials. *Arch Intern Med*. 2003;163:669–676.
3. Amarenco P, Bogousslavsky J, Callahan AS, Goldstein L, Hennerici M, Sillsen H, Welch MA, Zivin J, SPARCL Investigators. Design and baseline characteristics of the Stroke Prevention by Aggressive Reduction in Cholesterol Levels (SPARCL) study. *Cerebrovasc Dis*. 2003;16:389–395.

The American Heart Association makes every effort to avoid any actual or potential conflicts of interest that may arise as a result of an outside relationship or a personal, professional, or business interest of a member of the writing panel. Specifically, all members of the writing group are required to complete and submit a Disclosure Questionnaire showing all such relationships that might be perceived as real or potential conflicts of interest.

This statement was approved by the American Heart Association Science Advisory and Coordinating Committee on January 12, 2004. To purchase reprints call 410-528-4121, fax 410-528-4264, or e-mail kgray@lww.com. Ask for reprint No. 71-0282. To make photocopies for personal or educational use, call the Copyright Clearance Center, 978-750-8400.

(*Stroke* 2002;35:1023.)

© 2004 American Heart Association, Inc.

Stroke is available at <http://www.strokeaha.org>

DOI: 10.1161/01.STR.0000122762.96972.DD